

Table 28: *In vitro* antibacterial activities of crude extracts

Test Samples		Diameter/zones of inhibition (mm)						
		<i>S. aureus</i>	<i>S. faecalis</i>	<i>B. anthracis</i>	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>S. typhimurium</i>	<i>P. aeruginosa</i>
<i>L. eriocalyx</i> (Stem)	<i>n</i> -hexane	ND						
	DCM	6.2±0.1	6.1±0.1	7.4±0.2	7.9±0.1	6.2±0.2	3.4±0.1	5.8±0.1
	MeOH	5.3±0.2	5.0±0.4	5.3±0.4	5.1±0.2	6.3±0.3	8.4±0.3	5.4±0.4
<i>A. ovalifolius</i> (Root)	<i>n</i> -hexane	7.2±0.01	5.5±0.01	ND	ND	5.6±0.1	ND	ND
	EtOAc	5.0±0.4	5.2±0.04	5.5±0.06	5.6±0.01	5.6±0.51	5.0±0.08	ND
	DCM	15.2±0.1	6.0±0.1	4.6±0.1	10.8±0.1	10.3±0.1	5.0±0.1	5.1±0.1
	MeOH	12.3±0.1	6.2±0.1	5.6±0.1	8.1±0.1	8.0±0.2	13.3±0.1	6.1±0.2
<i>E. abyssinica</i> (Leaves)	<i>n</i> -Hexane	5.2±0.01	5.5±0.01	5.6±0.01	5.2±0.0	5.6±0.01	ND	ND
	EtOAc	5.2±0.07	5.5±0.01	5.6±0.01	ND	5.0±0.01	ND	ND
	DCM	15.3±0.1	7.0±0.1	4.6±0.1	10.8±0.1	10.3±0.1	5.0±0.1	4.1±0.1
	MeOH	11.3±0.1	6.2±0.1	5.6±0.1	8.1±0.1	5.0±0.2	5.3±0.1	5.1±0.2
	Am (20 µg/mL)	19.5±0.1	19.3±0.3	16.7±0.1	18.5±0.1	19.9±0.2	19.6±0.0	17.7±0.3

Key: *S. aureus*, (ATCC 25922), *S. faecalis* (ATCC 25925), *K. pneumoniae* (ATCC 90028), *S. typhimurium* (ATCC 25927), *E. coli* (K 12), *P. aeruginosa* (ATCC 25923), *B. anthracis* (QST 713), Am = Amoxycillin, ND = Not Detected

* Values are means ± SD of three replicates recorded at a concentration of 1000 µg/mL

Activity scale: (> 17: Highly active; 11-16: intermediate; 7-10: weak; <6: resistant) (Singh *et al.*, 2002, McChesney *et al.* 1991).